

FTUV/97-03
 IFIC/97-03
 January 1997

QCD TESTS FROM TAU DECAYS *

A. PICH

*Departament de Física Teòrica, IFIC, CSIC — Universitat de València,
 Dr. Moliner 50, E-46100 Burjassot, València, Spain*

The total τ hadronic width can be accurately calculated using analyticity and the operator product expansion. The theoretical analysis of this observable is updated to include all available perturbative and non-perturbative corrections. The experimental determination of $\alpha_s(M_\tau^2)$ and its actual uncertainties are discussed.

1 Introduction

The inclusive character of the total τ hadronic width renders possible an accurate calculation of the ratio¹⁻⁹ $[(\gamma)$ represents additional photons or lepton pairs]

$$R_\tau \equiv \frac{\Gamma[\tau^- \rightarrow \nu_\tau \text{ hadrons}(\gamma)]}{\Gamma[\tau^- \rightarrow \nu_\tau e^- \bar{\nu}_e(\gamma)]}, \quad (1)$$

using standard field theoretic methods. If strong and electroweak radiative corrections are ignored and if the masses of final-state particles are neglected, the universality of the W coupling to the fermionic charged currents implies

$$R_\tau \simeq N_c (|V_{ud}|^2 + |V_{us}|^2) \simeq 3, \quad (2)$$

which compares quite well with the experimental average¹⁰ $R_\tau = 3.649 \pm 0.014$. This provides strong evidence for the colour degree of freedom N_c .

The QCD dynamics is able to account quantitatively for the 20% difference between the naïve prediction (2) and the measured value of R_τ . Moreover, the uncertainties in the theoretical calculation of R_τ are quite small. The value of R_τ can then be accurately predicted as a function of $\alpha_s(M_\tau^2)$. Alternatively, measurements of inclusive τ decay rates can be used to determine the value of the QCD running coupling $\alpha_s(M_\tau^2)$ at the scale of the τ mass. In fact, τ decay is probably the lowest-energy process from which the running coupling constant can be extracted cleanly, without hopeless complications from non-perturbative effects. The τ mass,¹⁰ $M_\tau = (1777.00^{+0.30}_{-0.27})$ MeV, lies fortuitously in a compromise region where the coupling constant $\alpha_s(M_\tau^2)$ is large enough that R_τ is sensitive to its value, yet still small enough that the perturbative

*Invited talk at the 20th Johns Hopkins Workshop: *Non Perturbative Particle Theory & Experimental Tests* (Heidelberg, 27–29 June 1996)

expansion still converges well. Moreover, the non-perturbative contributions to the total τ hadronic width are very small.

It is the inclusive nature of the total semihadronic decay rate that makes a rigorous theoretical calculation of R_τ possible. The only separate contributions to R_τ that can be calculated are those associated with specific quark currents. We can calculate the non-strange and strange contributions to R_τ , and resolve these further into vector and axial-vector contributions. Since strange decays cannot be resolved experimentally into vector and axial-vector contributions, we will decompose our predictions for R_τ into only three categories:

$$R_\tau = R_{\tau,V} + R_{\tau,A} + R_{\tau,S}. \quad (3)$$

Non-strange semihadronic decays of the τ are resolved experimentally into vector ($R_{\tau,V}$) and axial-vector ($R_{\tau,A}$) contributions according to whether the hadronic final state includes an even or odd number of pions. Strange decays ($R_{\tau,S}$) are of course identified by the presence of an odd number of kaons in the final state. The naïve predictions for these three ratios are $R_{\tau,V} \simeq R_{\tau,A} \simeq (N_c/2)|V_{ud}|^2$ and $R_{\tau,S} \simeq N_c|V_{us}|^2$, which add up to (2).

2 Theoretical Framework

The theoretical analysis of R_τ involves the two-point correlation functions for the vector $V_{ij}^\mu = \bar{\psi}_j \gamma^\mu \psi_i$ and axial-vector $A_{ij}^\mu = \bar{\psi}_j \gamma^\mu \gamma_5 \psi_i$ colour-singlet quark currents ($i, j = u, d, s$):

$$\Pi_{ij,V}^{\mu\nu}(q) \equiv i \int d^4x e^{iqx} \langle 0 | T(V_{ij}^\mu(x) V_{ij}^\nu(0)^\dagger) | 0 \rangle, \quad (4)$$

$$\Pi_{ij,A}^{\mu\nu}(q) \equiv i \int d^4x e^{iqx} \langle 0 | T(A_{ij}^\mu(x) A_{ij}^\nu(0)^\dagger) | 0 \rangle. \quad (5)$$

They have the Lorentz decompositions

$$\Pi_{ij,V/A}^{\mu\nu}(q) = (-g^{\mu\nu} q^2 + q^\mu q^\nu) \Pi_{ij,V/A}^{(1)}(q^2) + q^\mu q^\nu \Pi_{ij,V/A}^{(0)}(q^2), \quad (6)$$

where the superscript (J) denotes the angular momentum $J = 1$ or $J = 0$ in the hadronic rest frame.

The imaginary parts of the two-point functions $\Pi_{ij,V/A}^{(J)}(q^2)$ are proportional to the spectral functions for hadrons with the corresponding quantum numbers. The semihadronic decay rate of the τ can be written as an integral of these spectral functions over the invariant mass s of the final-state hadrons:

$$R_\tau = 12\pi \int_0^{M_\tau^2} \frac{ds}{M_\tau^2} \left(1 - \frac{s}{M_\tau^2}\right)^2 \left[\left(1 + 2\frac{s}{M_\tau^2}\right) \text{Im}\Pi^{(1)}(s) + \text{Im}\Pi^{(0)}(s) \right]. \quad (7)$$




Figure 1: Integration contour in the complex s plane, used to obtain Eq. (9)

The appropriate combinations of correlators are

$$\Pi^{(J)}(s) \equiv |V_{ud}|^2 \left(\Pi_{ud,V}^{(J)}(s) + \Pi_{ud,A}^{(J)}(s) \right) + |V_{us}|^2 \left(\Pi_{us,V}^{(J)}(s) + \Pi_{us,A}^{(J)}(s) \right). \quad (8)$$

The contributions coming from the first two terms correspond to $R_{\tau,V}$ and $R_{\tau,A}$ respectively, while $R_{\tau,S}$ contains the remaining Cabibbo-suppressed contributions.

Since the hadronic spectral functions are sensitive to the non-perturbative effects of QCD that bind quarks into hadrons, the integrand in Eq. (7) cannot be calculated at present from QCD. Nevertheless the integral itself can be calculated systematically by exploiting the analytic properties of the correlators $\Pi^{(J)}(s)$. They are analytic functions of s except along the positive real s axis, where their imaginary parts have discontinuities. The integral (7) can therefore be expressed as a contour integral in the complex s plane running counter-clockwise around the circle $|s| = M_\tau^2$:

$$R_\tau = 6\pi i \oint_{|s|=M_\tau^2} \frac{ds}{M_\tau^2} \left(1 - \frac{s}{M_\tau^2} \right)^2 \left[\left(1 + 2 \frac{s}{M_\tau^2} \right) \Pi^{(0+1)}(s) - 2 \frac{s}{M_\tau^2} \Pi^{(0)}(s) \right]. \quad (9)$$

The advantage of expression (9) over (7) is that it requires the correlators only for complex s of order M_τ^2 , which is significantly larger than the scale associated with non-perturbative effects in QCD. The short-distance Operator Product Expansion (OPE) can therefore be used to organize the perturba-

tive and non-perturbative contributions to the correlators into a systematic expansion¹¹ in powers of $1/s$,

$$\Pi^{(J)}(s) = \sum_{D=2n} \sum_{\dim \mathcal{O}=D} \frac{\mathcal{C}^{(J)}(s, \mu) \langle \mathcal{O}(\mu) \rangle}{(-s)^{D/2}}, \quad (10)$$

where the inner sum is over local gauge-invariant scalar operators of dimension $D = 0, 2, 4, \dots$. The possible uncertainties associated with the use of the OPE near the time-like axis are absent in this case, because the integrand in Eq. (9) includes a factor $(1 - s/M_\tau^2)^2$, which provides a double zero at $s = M_\tau^2$, effectively suppressing the contribution from the region near the branch cut. The parameter μ in Eq. (10) is an arbitrary factorization scale, which separates long-distance non-perturbative effects, which are absorbed into the vacuum matrix elements $\langle \mathcal{O}(\mu) \rangle$, from short-distance effects, which belong in the Wilson coefficients $\mathcal{C}^{(J)}(s, \mu)$. The $D = 0$ term (unit operator) corresponds to the pure perturbative contributions, neglecting quark masses. The leading quark-mass corrections generate the $D = 2$ term. The first dynamical operators involving non-perturbative physics appear at $D = 4$. Inserting the functions (10) into (9) and evaluating the contour integral, R_τ can be expressed as an expansion in powers of $1/M_\tau^2$, with coefficients that depend only logarithmically on M_τ .

It is convenient to express the corrections to R_τ from dimension- D operators in terms of the fractional corrections $\delta_{ij,V/A}^{(D)}$ to the naïve contribution from the current with quantum numbers ij, V or ij, A :

$$\begin{aligned} R_{\tau,V/A} &= \frac{3}{2} |V_{ud}|^2 S_{EW} \left(1 + \delta'_{EW} + \sum_{D=2n} \delta_{ud,V/A}^{(D)} \right), \\ R_{\tau,S} &= 3 |V_{us}|^2 S_{EW} \left(1 + \delta'_{EW} + \sum_{D=2n} \delta_{us}^{(D)} \right). \end{aligned} \quad (11)$$

$\delta_{ij}^{(D)} = (\delta_{ij,V}^{(D)} + \delta_{ij,A}^{(D)})/2$ is the average of the vector and axial-vector corrections. The dimension-0 contribution is the purely perturbative correction neglecting quark masses, which is the same for all the components of R_τ : $\delta_{ij,V/A}^{(0)} = \delta^{(0)}$. The factors¹²

$$S_{EW} = \left(\frac{\alpha(m_b^2)}{\alpha(M_\tau^2)} \right)^{\frac{9}{19}} \left(\frac{\alpha(M_W^2)}{\alpha(m_b^2)} \right)^{\frac{9}{20}} \left(\frac{\alpha(M_Z^2)}{\alpha(M_W^2)} \right)^{\frac{36}{17}} = 1.0194 \quad (12)$$

and¹³

$$\delta'_{EW} = \frac{5}{12} \frac{\alpha(M_\tau^2)}{\pi} \simeq 0.0010, \quad (13)$$

contain the known electroweak corrections.

Adding the three terms, the total ratio R_τ is

$$R_\tau = 3 (|V_{ud}|^2 + |V_{us}|^2) S_{EW} \left\{ 1 + \delta'_{EW} + \delta^{(0)} + \sum_{D=2,4,\dots} \left(\cos^2 \theta_C \delta_{ud}^{(D)} + \sin^2 \theta_C \delta_{us}^{(D)} \right) \right\}, \quad (14)$$

where $\sin^2 \theta_C \equiv |V_{us}|^2 / (|V_{ud}|^2 + |V_{us}|^2)$.

3 Perturbative Corrections

In the chiral limit ($m_u = m_d = m_s = 0$), the vector and axial-vector currents are conserved. This implies $s\Pi^{(0)}(s) = 0$; therefore, only the correlator $\Pi^{(0+1)}(s)$ contributes to Eq. (9). Owing^{14,15} to the chiral invariance of massless QCD, $\Pi_{ij,V}^{(0+1)}(s) = \Pi_{ij,A}^{(0+1)}(s) \equiv \Pi(s)$ ($i \neq j$) at any finite order in α_s .

The result is more conveniently expressed in terms of the logarithmic derivative of the two-point correlation function of the vector (axial) current,

$$D(s) \equiv -s \frac{d}{ds} \Pi(s) = \frac{1}{4\pi^2} \sum_{n=0} K_n \left(\frac{\alpha_s(s)}{\pi} \right)^n, \quad (15)$$

which satisfies an homogeneous renormalization-group equation. The K_n coefficients are known^{16–20} to order α_s^3 . For $n_f = 3$ quark flavours, one has:

$$\begin{aligned} K_0 &= K_1 = 1, \\ K_2 &= \frac{299}{24} - 9\zeta(3) = 1.63982, \\ K_3(\overline{MS}) &= \frac{58057}{288} - \frac{779}{4}\zeta(3) + \frac{75}{2}\zeta(5) = 6.37101. \end{aligned} \quad (16)$$

The perturbative component of R_τ is given by

$$\frac{R_\tau^{\text{pert}}}{(|V_{ud}|^2 + |V_{us}|^2) S_{EW}} \equiv 3 \{1 + \delta^{(0)}\} = 3 \sum_{n=0} K_n A^{(n)}(\alpha_s), \quad (17)$$

where the functions⁵

$$A^{(n)}(\alpha_s) = \frac{1}{2\pi i} \oint_{|s|=M_\tau^2} \frac{ds}{s} \left(\frac{\alpha_s(-s)}{\pi} \right)^n \left(1 - 2 \frac{s}{M_\tau^2} + 2 \frac{s^3}{M_\tau^6} - \frac{s^4}{M_\tau^8} \right) \quad (18)$$

are contour integrals in the complex plane, which only depend on $\alpha_s(M_\tau^2)$.

The running coupling $\alpha_s(-s)$ in Eq. (18) can be expanded in powers of $\alpha_s(M_\tau^2)$, with coefficients that are polynomials in $\log(-s/M_\tau^2)$. The perturbative expansion of $\delta^{(0)}$ in powers of $a_\tau \equiv \alpha_s(M_\tau^2)/\pi$ then takes the form:¹⁻⁷

$$\begin{aligned}\delta^{(0)} &= \sum_{n=1} (K_n + g_n) a_\tau^n, \\ &= a_\tau + \left(K_2 - \frac{19}{24}\beta_1\right) a_\tau^2 \\ &\quad + \left(K_3 - \frac{19}{12}K_2\beta_1 - \frac{19}{24}\beta_2 + \frac{265 - 24\pi^2}{288}\beta_1^2\right) a_\tau^3 + \mathcal{O}(a_\tau^4) \\ &= a_\tau + 5.2023 a_\tau^2 + 26.366 a_\tau^3 + \mathcal{O}(a_\tau^4).\end{aligned}\tag{19}$$

The complex integration along the circle $|s| = M_\tau^2$ generates the g_n coefficients, which depend on $K_{m < n}$ and on $\beta_{m < n}$, where

$$\beta_1 = -9/2, \quad \beta_2 = -8, \quad \beta_3(\overline{MS}) = -3863/192, \quad \dots \tag{20}$$

are the coefficients of the QCD β function for $n_f = 3$. One observes⁵ that the g_n contributions are larger than the original K_n coefficients ($g_2 = 3.5625$, $g_3 = 19.9949$). For instance, the bold-guess value $K_4 \sim K_3(K_3/K_2) \approx 25$ is to be compared with $g_4 = 78.0029$. These large corrections give rise to a sizeable renormalization scale dependence^{5,21} of the truncated $\mathcal{O}(a_\tau^3)$ result. The reason of such uncomfortably large contributions⁵ stems from the long running along the circle $s = M_\tau^2 \exp(i\phi)$ ($\phi \in [0, 2\pi]$) in Eq. (18). When the running coupling $\alpha_s(-s)$ is expanded in powers of $\alpha_s(M_\tau^2)$, one gets imaginary logarithms, $\log(-s/M_\tau^2) = i(\phi - \pi)$, which are large in some parts of the integration range. The radius of convergence of this expansion is actually quite small. A numerical analysis of the series⁵ shows that, at the three-loop level, an upper estimate for the convergence radius is $a_{\tau, \text{conv}} < 0.11$.

Note, however, that there is no deep reason to stop the $A^{(n)}(\alpha_s)$ integral expansions at $\mathcal{O}(\alpha_s^3)$. One can calculate the $A^{(n)}(\alpha_s)$ expansion to all orders in α_s , apart from the unknown $\beta_{n>3}$ contributions, which are likely to be negligible. Even for a_τ larger than the radius of convergence $a_{\tau, \text{conv}}$, the integrals $A^{(n)}(\alpha_s)$ are well-defined functions that can be numerically computed, by using in Eq. (18) the exact solution for $\alpha_s(-s)$ obtained from the renormalization-group β -function equation. Thus a more appropriate approach^{5,22} is to use a K_n expansion of R_τ^{pert} as in Eq. (17), and to fully keep the known three-loop calculation of the functions $A^{(n)}(\alpha_s)$. The perturbative uncertainties are then reduced to the corrections coming from the unknown $\beta_{n>3}$ and $K_{n>3}$ contributions, since the g_n contributions are properly resummed to all orders. To

appreciate the size of the effect, Table 1 gives the exact results⁵ for $A^{(n)}(\alpha_s)$ ($n = 1, 2, 3$) obtained at the one-, two- and three-loop approximations (i.e. $\beta_{n>1} = 0$, $\beta_{n>2} = 0$, and $\beta_{n>3} = 0$, respectively), together with the final value of $\delta^{(0)}$, for $a_\tau = 0.1$. For comparison, the numbers coming from the truncated expressions at order a_τ^3 are also given. Although the difference between the exact and truncated results represents a tiny 0.7% effect on R_τ , it produces a sizeable 4% shift on the value of $\delta^{(0)}$. The $\delta^{(0)}$ shift, which reflects into a corresponding shift in the experimental $\alpha_s(M_\tau^2)$ determination, depends strongly on the value of the coupling constant; for $a_\tau = 0.14$ the $\delta^{(0)}$ shift reaches the -20% level.

Table 1: Exact results for $A^{(n)}(\alpha_s)$ ($n = 1, 2, 3$) obtained at the one- ($\beta_{n>1} = 0$), two- ($\beta_{n>2} = 0$) and three-loop ($\beta_{n>3} = 0$) approximations, together with the final value of $\delta^{(0)}$, for $a_\tau = 0.1$. For comparison, the numbers coming from the truncated expressions at order a_τ^3 are also given.

Loops	$A^{(1)}$	$A^{(2)}$	$A^{(3)}$	$\delta^{(0)}$
1	0.13247	0.01570	0.00170	0.1690
2	0.13523	0.01575	0.00163	0.1714
3	0.13540	0.01565	0.00160	0.1712
$\mathcal{O}(\alpha_s^3)$	0.14394	0.01713	0.00100	0.1784

Notice that the difference between using the one- or two-loop approximation to the β function is already quite small (1.4% effect on $\delta^{(0)}$), while the change induced by the three-loop corrections is completely negligible (0.1%). Therefore (unless the β function has some unexpected pathological behaviour at higher orders), the error induced by the truncation of the β function at third order should be smaller than 0.1% and therefore can be safely neglected. The sensitivity on the choice of renormalization scale and renormalization scheme is also very small.^{5,23}

The dominant perturbative uncertainties come from the unknown higher-order coefficients $K_{n>3}$. The $\mathcal{O}(\alpha_s^4)$ contribution has been estimated²⁴ using scheme-invariant methods, namely the principle of minimal sensitivity²⁵ and the effective charge approach,²⁶ with the result:²⁴

$$K_4^{\text{est}} = 27.5. \quad (21)$$

This number is very close to the naïve guess⁵ $K_4 \sim (K_3/K_2)K_3 \approx 25$. A similar estimate, $K_4^{\text{NNA}} = 24.8$, is obtained^{27–33} in the limit of a large number of quark flavours, using the so-called *naïve non-abelianization* prescription³⁴

Table 2: $\delta^{(0)}$ for different values of $\alpha_s(M_\tau^2)$.

$\alpha_s(M_\tau^2)$	$\delta^{(0)}$		$\Delta(\delta^{(0)})$
	$K_4 = 0$	$K_4 = 27.5$	
0.30	0.161	0.164	± 0.006
0.31	0.168	0.172	± 0.007
0.32	0.176	0.180	± 0.008
0.33	0.183	0.188	± 0.008
0.34	0.191	0.196	± 0.009
0.35	0.198	0.203	± 0.010
0.36	0.205	0.211	± 0.010
0.37	0.213	0.219	± 0.011
0.38	0.220	0.226	± 0.012
0.39	0.227	0.234	± 0.012
0.40	0.234	0.241	± 0.013

($n_f \rightarrow 3\beta_1 = n_f - \frac{33}{2} = -\frac{27}{2}$). From a fit to the experimental τ data, the value $K_4^{\text{fit}} = 29 \pm 5$ has been also quoted.³⁵

Using the estimate (21), the $\mathcal{O}(\alpha_s^4)$ correction amounts to a 0.004 increase of $\delta^{(0)}$ for $a_\tau = 0.1$. The resulting perturbative contribution $\delta^{(0)}$, obtained through Eqs. (17) and (18), is given in Table 2 for different values of the strong coupling constant $\alpha_s(M_\tau^2)$. In order to be conservative, and to account for all possible sources of perturbative uncertainties, we have used^{6,7}

$$\Delta(\delta^{(0)}) = \pm 50 A^{(4)}(\alpha_s), \quad (22)$$

as an estimate of the theoretical error on $\delta^{(0)}$. Note that, for the relevant values of α_s , this is of the same size as $K_3 A^{(3)}(\alpha_s)$; thus, this error estimate is conservative enough to apply⁷ in the worst possible scenario, where the onset of the asymptotic behaviour of the perturbative series were already reached for $n = 3, 4$.

There have been attempts^{29–33} to improve the perturbative prediction by performing an all-order summation a certain class of higher-order corrections (the so-called ultraviolet renormalon chains). This can be accomplished using exact large- n_f results and applying the *naive non-abelianization* prescription.³⁴ Unfortunately, the naive resummation turns out to be renormalization-scheme dependent beyond one loop.^{30,36} More recently, a renormalization-scheme-invariant summation has been presented.³⁷ The final effect of the

higher-order corrections (beyond K_4) turns out to be small.^a

4 Power Corrections

The $1/M_\tau^2$ contributions $\delta_{ij}^{(2)}$ to the ratio R_τ are simply the leading quark-mass corrections to the perturbative QCD result of the previous section. These contributions are known^{4,38} to order α_s^2 . Quark-mass corrections are certainly tiny for the up and down quarks ($\delta_{ud}^{(2)} \sim -0.08\%$), but the correction from the strange-quark mass is important for strange decays:^{4,38} ($m_u = m_d = 0$)

$$\delta_{us,V/A}^{(2)} = -8 \frac{\overline{m}_s^2}{M_\tau^2} \left\{ 1 + \frac{16}{3} a_\tau + 11.03 a_\tau^2 \right\}, \quad (23)$$

where $\overline{m}_s \equiv m_s(M_\tau^2)$ is the running mass of the strange quark evaluated at the scale M_τ . For $\alpha_s(M_\tau^2) = 0.35$, $\delta_{us}^{(2)} \approx -20\%$; nevertheless, because of the $\sin^2 \theta_C$ suppression, the effect on the total ratio R_τ is only $-(1.0 \pm 0.2)\%$.

Since the τ mass is a quite low energy scale, we should worry about possible non-perturbative effects. In the framework of the OPE, the long-distance dynamics is absorbed into the vacuum matrix elements $\langle \mathcal{O}(\mu) \rangle$, which are (at present) quantities to be fixed phenomenologically. If the logarithmic dependence of the Wilson coefficients $\mathcal{C}^{(J)}(s, \mu)$ on s is neglected (this is an effect of order α_s^2), the contour integrals can be evaluated trivially using Cauchy's residue theorem, and are non-zero only for $D = 2, 4, 6$ and 8 . The corrections simplify even further if we also take the chiral limit ($m_u = m_d = m_s = 0$). The dimension-2 corrections then vanish because there are no operators of dimension 2. In the chiral limit, $s\Pi^{(0)}(s) = 0$; thus only the $\Pi^{(0+1)}(s)$ term in Eq. (9) contributes to R_τ . The form of the kinematical factor multiplying $\Pi^{(0+1)}(s)$ in Eq. (9) is such that, when the s -dependence of the Wilson coefficients is ignored, only the $D = 6$ and $D = 8$ contributions survive the integration. The power corrections to R_τ then reduce to¹⁻⁴

$$\begin{aligned} \delta_{ij,V/A}^{(6)} &\simeq -24\pi^2 \frac{\left[\sum \mathcal{C}_{ij,V/A}^{(0+1)} \langle \mathcal{O} \rangle \right]^{(D=6)}}{M_\tau^6}, \\ \delta_{ij,V/A}^{(8)} &\simeq -16\pi^2 \frac{\left[\sum \mathcal{C}_{ij,V/A}^{(0+1)} \langle \mathcal{O} \rangle \right]^{(D=8)}}{M_\tau^8}, \end{aligned} \quad (24)$$

and $\delta_{ij,V/A}^{(2n)} \simeq 0$ for $2n \neq 6, 8$.

^a The present results do not include yet the resummation of the g_n running corrections through the $A^{(n)}(\alpha_s)$ functions.

When the logarithmic dependence of the Wilson coefficients on s is taken into account, operators of dimensions other than 6 and 8 do contribute, but they are suppressed by two powers of $\alpha_s(M_\tau^2)$. The largest power corrections to R_τ then come from dimension-6 operators, which have no such suppression. Their size was first estimated in Ref. 4, using published phenomenological fits to different sets of data:

$$\delta_{ij}^{(6)} \approx -\frac{2}{7}\delta_{ij,V}^{(6)} \approx \frac{2}{11}\delta_{ij,A}^{(6)} \approx -(0.7 \pm 0.4)\%. \quad (25)$$

These power corrections are numerically very small, which is due to the fact that they fall off like the sixth power of $1/M_\tau$. Moreover, there is a large cancellation between the vector and axial-vector contributions to the total hadronic width (the operator with the largest Wilson coefficient contributes with opposite signs to the vector and axial-vector correlators, due to the γ_5 flip). Thus, the non-perturbative corrections to R_τ are smaller than the corresponding contributions to $R_{\tau,V/A}$. A more detailed study of non-perturbative corrections, including the very small $D = 4$ contributions proportional to quark masses or to $\alpha_s(M_\tau^2)^2$, can be found in Ref. 4.

The estimate (25) introduces a small uncertainty in the R_τ predictions, since the actual evaluation of the non-perturbative contributions involves a mixture of experimental measurements and theoretical considerations, which are model-dependent to some extent. It is better to directly measure those contributions from the τ -decay data themselves. This information can be extracted³⁹ from the invariant-mass distribution of the final hadrons in τ decay.

Although the distributions themselves cannot be predicted at present, certain weighted integrals of the hadronic spectral functions can be calculated in the same way as R_τ . The analyticity properties of $\Pi_{ij,V/A}^{(J)}$ imply:^{39, 40}

$$\int_0^{s_0} ds W(s) \text{Im}\Pi_{ij,V/A}^{(J)} = \frac{i}{2} \oint_{|s|=s_0} ds W(s) \Pi_{ij,V/A}^{(J)}, \quad (26)$$

with $W(s)$ an arbitrary weight function without singularities in the region $|s| \leq s_0$. Generally speaking, the accuracy of the theoretical predictions can be much worse than the one of R_τ , because non-perturbative effects are not necessarily suppressed. In fact, choosing an appropriate weight function, non-perturbative effects can even be made to dominate the final result. But this is precisely what makes these integrals interesting: they can be used to measure the parameters characterizing the non-perturbative dynamics.

To perform an experimental analysis, it is convenient to use moments of

the directly measured invariant-mass distribution⁴⁰ ($k, l \geq 0$)

$$R_\tau^{kl}(s_0) \equiv \int_0^{s_0} ds \left(1 - \frac{s}{s_0}\right)^k \left(\frac{s}{M_\tau^2}\right)^l \frac{dR_\tau}{ds}. \quad (27)$$

The factor $(1 - s/s_0)^k$ supplements $(1 - s/M_\tau^2)^2$ for $s_0 \neq M_\tau^2$, in order to squeeze the integrand at the crossing of the positive real axis and, therefore, improves the reliability of the OPE analysis; moreover, for $s_0 = M_\tau^2$ it reduces the contribution from the tail of the distribution, which is badly defined experimentally. A combined fit of different $R_\tau^{kl}(s_0)$ moments results in experimental values for $\alpha_s(M_\tau^2)$ and for the coefficients of the inverse power corrections in the OPE. $R_\tau^{00}(M_\tau^2) = R_\tau$ uses the overall normalization of the hadronic distribution, while the ratios $D_\tau^{kl}(M_\tau^2) = R_\tau^{kl}(M_\tau^2)/R_\tau$ are based on the shape of the s distribution and are more dependent on non-perturbative effects.⁴⁰

The predicted suppression¹⁻⁴ of the non-perturbative corrections has been confirmed by ALEPH^{41,42} and CLEO,⁴³ using the moments (0,0), (1,0), (1,1), (1,2) and (1,3). The most recent ALEPH analysis⁴⁴ gives:

$$\delta_{\text{NP}} \equiv \sum_{D \geq 4} \left(\cos^2 \theta_C \delta_{ud}^{(D)} + \sin^2 \theta_C \delta_{us}^{(D)} \right) = -(0.5 \pm 0.7)\%, \quad (28)$$

in agreement with (25).

5 Phenomenology

The QCD prediction for R_τ is then completely dominated by the perturbative contribution $\delta^{(0)}$; non-perturbative effects being of the order of the perturbative uncertainties from uncalculated higher-order corrections.⁷⁻⁹ Furthermore, as shown in Table 2, the result turns out to be very sensitive to the value of $\alpha_s(M_\tau^2)$, allowing for an accurate determination of the fundamental QCD coupling.

The experimental value for R_τ can be obtained from the leptonic branching fractions or from the τ lifetime. The average of those determinations¹⁰

$$R_\tau = 3.649 \pm 0.014, \quad (29)$$

corresponds to

$$\alpha_s(M_\tau^2) = 0.355 \pm 0.025. \quad (30)$$

Once the running coupling constant $\alpha_s(s)$ is determined at the scale M_τ , it can be evolved to higher energies using the renormalization group. The size of its error bar scales roughly as α_s^2 , and it therefore shrinks as the scale increases.




Figure 2: Constraints on $\delta_V^{(6)}$ and $\delta_A^{(6)}$ obtained from ALEPH data.⁴⁴ The ellipse depicts the combined fit. All results are still preliminary. (Taken from Ref. 44)

Thus a modest precision in the determination of α_s at low energies results in a very high precision in the coupling constant at high energies. After evolution up to the scale M_Z , the strong coupling constant in (30) decreases to^b

$$\alpha_s(M_Z^2) = 0.1225 \pm 0.0025, \quad (31)$$

in excellent agreement with the direct measurement^{46,47} at $\mu = M_Z$, $\alpha_s(M_Z^2) = 0.121 \pm 0.003$, and with a similar error bar. The comparison of these two determinations of α_s in two extreme energy regimes, M_τ and M_Z , provides a beautiful test of the predicted running of the QCD coupling.

With $\alpha_s(M_\tau^2)$ fixed to the value in Eq. (30), the same theoretical framework gives definite predictions^{4,7} for the semi-inclusive τ decay widths $R_{\tau,V}$, $R_{\tau,A}$ and $R_{\tau,S}$, in good agreement with the experimental measurements.^{44,48} The separate analysis of the vector and axial-vector contributions allows to investigate the associated non-perturbative corrections ($R_{\tau,V} - R_{\tau,A}$ is a pure non-perturbative quantity). Figure 2 shows⁴⁴ the (preliminary) constraints on $\delta_V^{(6)}$ and $\delta_A^{(6)}$ obtained from the most recent ALEPH analyses.^{41,44,45} A clear improvement over previous phenomenological determinations^{4,49} is apparent.

^b From a combined analysis of τ data, ALEPH quotes:⁴⁴ $\alpha_s(M_Z^2) = 0.1225 \pm 0.0006_{\text{exp}} \pm 0.0015_{\text{th}} \pm 0.0010_{\text{evol}}$.




Figure 3: Values of $\alpha_s(s_0)$ extracted⁵⁰ from the $R_\tau^{00}(s_0)$ data.^{42, 43} The dashed line shows the three-loop QCD prediction for the running coupling constant. (Taken from Ref. 50)

The Cabibbo-suppressed width $R_{\tau,S}$ is very sensitive to the value of the strange quark mass,⁴ providing a direct and clean way of measuring m_s . A very preliminary value, $m_s(M_\tau^2) = (212^{+30+1}_{-35-5})$ MeV, has been already presented at the last tau workshop.⁴⁸

Using the measured invariant-mass distribution of the final hadrons, it is possible to evaluate the integral $R_\tau^{00}(s_0)$, with an arbitrary upper limit of integration $s_0 \leq M_\tau^2$. The experimental s_0 dependence agrees well with the theoretical predictions⁴⁰ up to rather low values⁵⁰ of s_0 (> 0.7 GeV²). Equivalently, from the measured^{42,43} $R_\tau^{00}(s_0)$ distribution one obtains $\alpha_s(s_0)$ as a function of the scale s_0 . As shown⁵⁰ in Figure 3, the result exhibits an impressive agreement with the running predicted at three-loop order by QCD. It is important to realize⁵⁰ that the theoretical prediction for $R_\tau^{00}(s_0)$ does not contain inverse powers of s_0 (as long as the s -dependence of the Wilson coefficients is ignored). The power corrections are suppressed by powers of $1/M_\tau^2$; thus, they do not drive a break-down of the OPE. This could explain the surprisingly good agreement with the data for $s_0 \lesssim 1$ GeV².

A similar test was performed before⁵¹ for $R_{\tau,V}$, using the vector spectral function measured in $e^+e^- \rightarrow \text{hadrons}$, and varying the value of the tau mass. This allows to study the behaviour of the OPE at lower scales. The theoretical predictions for $R_{\tau,V}$ as function of M_τ^2 agree⁵¹ well with the data for $M_\tau > 1.2$ GeV. Below this value, higher-order inverse power corrections become very important and eventually generate the expected break-down of the expansion in powers of $1/M_\tau^2$.

6 Summary

Because of its inclusive nature, the total hadronic width of the τ can be rigorously computed within QCD. One only needs to study two-point correlation functions for the vector and axial-vector currents. As shown in Eq. (9), this information is only needed in the complex plane, away from the time-like axis; the dangerous region near the physical cut does not contribute at all to the result, because of the phase-space factor $(1 - s/M_\tau^2)^2$. The uncertainties of the theoretical predictions are then quite small.

The ratio R_τ is very sensitive to the value of the strong coupling, and therefore can be used² to measure $\alpha_s(M_\tau^2)$. This observation has triggered an ongoing effort to improve the knowledge of R_τ from both the experimental and the theoretical sides. The fact that M_τ is a quite low energy scale (i.e. that $\alpha_s(M_\tau^2)$ is big), but still large enough to allow a perturbative analysis, makes R_τ an ideal observable to determine the QCD coupling. Moreover, since the error of $\alpha_s(\mu^2)$ shrinks as μ increases, the good accuracy of the R_τ determination of $\alpha_s(M_\tau^2)$ implies a very precise value of $\alpha_s(M_Z^2)$.

The theoretical analysis of R_τ has reached a very mature level. Many different sources of possible perturbative and non-perturbative contributions have been analyzed in detail. A very detailed study of the associated uncertainties has been given in Ref. 7. The final theoretical uncertainty is small and has been adequately taken into account in the final $\alpha_s(M_\tau^2)$ determination in Eq. (30).

The comparison of the theoretical predictions with the experimental data shows a successful and consistent picture. The $\alpha_s(M_\tau^2)$ determination is in excellent agreement with the measurements at the Z -mass scale, providing clear evidence of the running of α_s . Moreover, the analysis of the semi-inclusive components of the τ hadronic width, $R_{\tau,V}$, $R_{\tau,A}$ and $R_{\tau,S}$, and the invariant-mass distribution of the final decay products gives a further confirmation of the reliability of the theoretical framework, and allows to investigate other important QCD parameters such as the strange-quark mass or the non-perturbative vacuum condensates.

Acknowledgments

I would like to thank the organizers for this enjoyable meeting, and specially Matthias Jamin for his kind hospitality. I am also indebted to Andreas Höcker and Matthias Neubert for providing figures 2 and 3. This work has been supported in part by CICYT (Spain) under grant No. AEN-96-1718.

References

1. E. Braaten, *Phys. Rev. Lett.* **60** (1988) 1606; *Phys. Rev.* **D39** (1989) 1458.
2. S. Narison and A. Pich, *Phys. Lett.* **B211** (1988) 183.
3. A. Pich, *Hadronic Tau-Decays and QCD*, Proc. Workshop on Tau Lepton Physics (Orsay, 1990), eds. M. Davier and B. Jean-Marie (Ed. Frontières, Gif-sur-Yvette, 1991), p. 321.
4. E. Braaten, S. Narison and A. Pich, *Nucl. Phys.* **B373** (1992) 581.
5. F. Le Diberder and A. Pich, *Phys. Lett.* **B286** (1992) 147.
6. A. Pich, *QCD Predictions for the Tau Hadronic Width and Determination of $\alpha_s(m_\tau^2)$* , Proc. Second Workshop on Tau Lepton Physics (Ohio, 1992), ed. K.K. Gan (World Scientific, Singapore, 1993), p. 121.
7. A. Pich, *Nucl. Phys. B (Proc. Suppl.)* **39B,C** (1995) 326.
8. S. Narison, *Nucl. Phys. B (Proc. Suppl.)* **40** (1995) 47.
9. E. Braaten, *Perturbative QCD and Tau Decay*, Proc. Fourth Workshop on Tau Lepton Physics –TAU96– (Colorado, 16–19 September 1996), ed. J. Smith, *Nucl. Phys. B (Proc. Suppl.)* in press.
10. A. Pich, *Tau Lepton Physics: Theory Overview*, Proc. Fourth Workshop on Tau Lepton Physics –TAU96– (Colorado, 16–19 September 1996), ed. J. Smith, *Nucl. Phys. B (Proc. Suppl.)* in press [IFIC/96-96; hep-ph/9612308].
11. M.A. Shifman, A.L. Vainshtein and V.I. Zakharov, *Nucl. Phys.* **B147** (1979) 385; 448; 519.
12. W.J. Marciano and A. Sirlin, *Phys. Rev. Lett.* **61** (1988) 1815; **56** (1986) 22.
13. E. Braaten and C.S. Li, *Phys. Rev.* **D42** (1990) 3888.
14. T.L. Trueman, *Phys. Lett.* **88B** (1979) 331.
15. A.I. Antoniadis, *Phys. Lett.* **84B** (1979) 223.
16. K.G. Chetyrkin, A.L. Kataev and F.V. Tkachov, *Phys. Lett.* **85B** (1979) 277.
17. M. Dine and J. Sapiirstein, *Phys. Rev. Lett.* **43** (1979) 668.
18. W. Celmaster and R. Gonsalves, *Phys. Rev. Lett.* **44** (1980) 560.
19. S.G. Gorishny, A.L. Kataev and S.A. Larin, *Phys. Lett.* **B259** (1991) 144.
20. L.R. Surguladze and M.A. Samuel, *Phys. Rev. Lett.* **66** (1991) 560.
21. J. Chýla, A. Kataev and S. Larin, *Phys. Lett.* **B267** (1991) 269.
22. A.A. Pivovarov, *Z. Phys.* **C53** (1992) 461.
23. P.A. Raczka and A. Szymacha, *Z. Phys.* **C70** (1996) 125.
24. A.L. Kataev and V.V. Starshenko, *Mod. Phys. Lett.* **A10** (1995) 235.

25. P.M. Stevenson, *Phys. Rev.* **D23** (1981) 2916.
26. G. Grunberg, *Phys. Lett.* **B221** (1980) 70; *Phys. Rev.* **D29** (1984) 2315.
27. D.J. Broadhurst, *Z. Phys.* **C58** (1993) 339.
28. D.J. Broadhurst and A.L. Kataev, *Phys. Lett.* **B315** (1993) 179.
29. M. Beneke, *Nucl. Phys.* **B405** (1993) 424; *Phys. Lett.* **B307** (1993) 154.
30. C.N. Lowett–Turner and C.G. Maxwell, *Nucl. Phys.* **B452** (1995) 188; **432** (1994) 147.
31. M. Beneke and V.M. Braun, *Phys. Lett.* **B348** (1995) 513.
32. P. Ball, M. Beneke and V.M. Braun, *Nucl. Phys.* **B452** (1995) 563.
33. M. Neubert, *Nucl. Phys.* **B463** (1996) 511.
34. D.J. Broadhurst and A.G. Grozin, *Phys. Rev.* **D52** (1995) 4082.
35. F. Le Diberder, *Nucl. Phys. B (Proc. Suppl.)* **39B,C** (1995) 318.
36. J. Chýla, *Phys. Lett.* **B356** (1995) 341.
37. C.J. Maxwell and D.G. Tonge, *RS-Invariant All-Orders Renormalon Resummations for some QCD Observables*, [hep-ph/9606392].
38. K.G. Chetyrkin and A. Kwiatkowski, *Z. Phys.* **C59** (1993) 525.
39. A. Pich, *QCD Tests From Tau Decay Data*, in Proc. Tau-Charm Factory Workshop (SLAC, 1989), ed. L.V. Beers, SLAC-Report-343 (1989), p. 416.
40. F. Le Diberder and A. Pich, *Phys. Lett.* **B289** (1992) 165.
41. D. Buskulic *et al* (ALEPH), *Phys. Lett.* **B307** (1993) 209.
42. L. Duflot, *Nucl. Phys. B (Proc. Suppl.)* **40** (1995) 37.
43. T. Coan *et al* (CLEO), *Phys. Lett.* **B356** (1995) 580.
44. A. Höcker, *Vector and Axial-Vector Spectral Functions and QCD*, Proc. Fourth Workshop on Tau Lepton Physics –TAU96– (Colorado, 16–19 September 1996), ed. J. Smith, *Nucl. Phys. B (Proc. Suppl.)* in press.
45. D. Buskulic *et al* (ALEPH), *Z. Phys.* **C70** (1996) 579.
46. M. Schmelling, *Status of the Strong Coupling Constant*, Proc. XXVIII Intern. Conf. on High Energy Physics (Warsaw, 25–31 July 1996) in press [hep-ex/9701002].
47. S. Bethke, *Experimental Tests of Asymptotic Freedom*, Proc. QCD’96 (Montpellier, 1996), ed. S. Narison [hep-ex/9609014].
48. M. Davier, *τ Decays into Strange Particles and QCD*, Proc. Fourth Workshop on Tau Lepton Physics –TAU96– (Colorado, 16–19 September 1996), ed. J. Smith, *Nucl. Phys. B (Proc. Suppl.)* in press.
49. S. Narison, *Phys. Lett.* **B361** (1995) 121.
50. M. Girone and M. Neubert, *Phys. Rev. Lett.* **76** (1996) 3061.
51. S. Narison and A. Pich, *Phys. Lett.* **B304** (1993) 359.